


Drug Information Sheet ("Kusuri-no-Shiori")

620003664

Revised: 02/2008

<p>Information is based on approvals granted in Japan. In some cases, the approved details may vary in countries other than Japan. In deciding to use a medicine, the risk (side effects) of taking the medicine must be weighed against the benefit (effects) it will do. The patient's cooperation is indispensable here.</p>	
<p>Brand name : LEUCON TABLETS 10mg Active ingredient : Adenine Dosage form : white tablet (: 6.4 mm, thickness: 3.0 mm) Printings on wrapping : Leucon, OH-277, ロイコン, ロイコン 10mg</p>	
<p>Effects of this medicine This medicine is taken up by RNA or DNA in bone marrow cells, is used for synthesis of nucleic acid, and consequently suppresses decrease of leucocytes and increases the leucocytes. It is usually used for the treatment of leukopenia induced by exposure to irradiation or drugs.</p>	
<p>Before using this medicine, tell your doctor/pharmacist</p> <ul style="list-style-type: none"> • If you ever experienced any allergic reaction (itch, rash, etc.) to any medicine. If you are a patient with gout or urolithiasis. • If you are pregnant or breastfeeding. • If you are using any other prescription or nonprescription (OTC) medicine. (It is important to reduce the risk of drug interactions.) 	
<p>Dosage regimen (proper use of this medicine)</p> <ul style="list-style-type: none"> • Your dosage regimen is: _____ :order of your doctor • General dosage regimen: For adults, take 2 to 6 tablets (20 to 60 mg of the active ingredient) daily, however, the dosage may be adjusted according to the age or symptoms. Strictly follow the instructions of your doctor/pharmacist. • Missed dose: Take a dose as soon as possible when you remembered that you missed a dose. However, if it is almost time for the next dose, skip the missed dose and continue your regular dosing schedule. DO NOT take double doses to make up for the missed dose. • Overdose: If you took too much of the medicine (more than ordered), check with your doctor/pharmacist. • Others: Do not stop taking the medicine without the instructions of your doctor. 	
<p>Precautions while using this medicine</p>	
<p>Side effects of this medicine Common side effects are reported as below. If any of them occurs, check with your doctor/pharmacist: rash, itch, etc. In rare cases, symptoms described below may be the sign of side effects indicated in brackets []. If they occur, stop taking your medicine and check with your doctor immediately.</p> <ul style="list-style-type: none"> • swelling accompanied by severe pain in the joint of the first toe, fever [hyperuricemia, gout] • low back pain, back pain, hematuria, decrease of urine output [urolithiasis, acute renal failure] <p>Other side effects not listed here may occur. Check with your doctor/pharmacist if you have any worrisome symptom.</p>	
<p>Storage and other information</p> <ul style="list-style-type: none"> • Keep the medicine out of the reach of children. Store it away from direct sunlight, heat and moisture. • Discard the remainder. Do not store them. 	
<p>For doctor use only: Day Month Year</p>	

For further information, ask your doctor/pharmacist. Japanese package insert information (for medical professionals) is available on the website of Pharmaceuticals and Medical Devices Agency.

SIE