

Drug Information Sheet("Kusuri-no-Shiori")

Internal

Revised: 06/2015

The information on this sheet is based on approvals granted by the Japanese regulatory authority. Approval details may vary by country. Medicines have adverse reactions (risks) as well as efficacies (benefits). It is important to minimize adverse reactions and maximize efficacy. To obtain a better therapeutic response, patients should understand their medication and cooperate with the treatment.

Brand name: LISINOPRIL TABLETS 10mg 「OHARA」

Active ingredient: Lisinopril hydrate

Dosage form: white butterfly-shaped tablet with split line, major axis: 9.3 mm, minor axis: 5.6 mm, thickness: 3.3 mm

Print on wrapping: リシノプリル「オーハラ」, 10mg, Lisinopril 10mg「OHARA」, リシノプリル 10mg「オーハラ」, リシノプリル, OH-77, 高血圧・慢性心不全のお薬です



Effects of this medicine

This medicine inhibits angiotensin-conversion enzyme to suppress production of angiotensin II. It subsequently dilates blood vessels and decreases blood pressure to reduce cardiac load.

It is usually used to treat hypertension and chronic heart failure (mild to moderate).

Before using this medicine, be sure to tell your doctor and pharmacist

- If you have previously experienced any allergic reactions (itch, rash, etc.) to any medicines.
If you have a history of angioedema.
If you are on apheresis or hemodialysis.
If you have renal artery stenosis, hyperkalemia, renal dysfunction or cerebrovascular disorder.
- If you are pregnant, possibly pregnant or breastfeeding.
- If you are taking any other medicinal products. (Some medicines may interact to enhance or diminish medicinal effects. Beware of over-the-counter medicines and dietary supplements as well as other prescription medicines.)

Dosing schedule (How to take this medicine)

- Your dosing schedule prescribed by your doctor is<< to be written by a healthcare professional>>
 - **Hypertension:** In general, for adults, take 1 to 2 tablets (10 to 20 mg of the active ingredient anhydride) at a time, once a day. The dosage may be adjusted according to your age and symptoms. For patient with severe hypertension or hypertension with renal disorder, the initial dosage may be started with 1/2 tablet (5 mg), once a day. In general, for children aged 6 years or over, take 0.07 mg/kg of the active ingredient anhydride at a time, once a day. The dosage may be adjusted according to your age and symptoms. However, the daily dosage should not exceed 20 mg. This preparation contains 10 mg of the active ingredient anhydride in a tablet.
 - **Chronic heart failure (mild to moderate):** This medicine is used in combination with base treatment such as digitalis products and diuretics. In general, for adults, take 1/2 to 1 tablet (5 to 10 mg of the active ingredient anhydride) at a time, once a day. The dosage may be adjusted according to your age and symptoms. For patient with renal disorder, the initial dosage may be started with 2.5 mg at a time, once a day.
- Strictly follow the instructions in any case.
- If you miss a dose, take the missed dose as soon as possible. However, if it is almost time for the next dose, skip the missed dose and continue your regular dosing schedule. You should never take two doses at one time.
 - If you accidentally take more than your prescribed dose, consult with your doctor or pharmacist.
 - Do not stop taking this medicine unless your doctor instructs you to do so.

Precautions while taking this medicine

- This medicine may cause dizziness or lightheadedness due to decreased blood pressure. Pay attention when performing dangerous operations such as working at heights or driving a car.

Possible adverse reactions to this medicine

The most commonly reported adverse reactions include dry cough, rash, itching and photosensitivity. If any of these symptoms occur, consult with your doctor or pharmacist.

The symptoms described below are rarely seen as initial symptoms of the adverse reactions indicated in brackets. If any of these symptoms occur, stop taking this medicine and see your doctor immediately.

- swelling of face/tongue/glottis/larynx with respiratory distress, abdominal pain, nausea [angioedema]
- decreased urine output, edema, thirst [acute renal failure]
- numbness in limbs/lips, muscular degeneration, paralysis in limbs [hyperkalemia]
- pain in upper abdomen/lower back, fever, nausea [pancreatitis]
- high fever more than 38°C, erythema/blister, erosion in lips/ocular mucosa [mucocutaneous ocular syndrome, toxic epidermal necrosis, pemphigus]

The above symptoms do not describe all the adverse reactions to this medicine. Consult with your

doctor or pharmacist if you notice any symptoms of concern other than those listed above.
Storage conditions and other information <ul style="list-style-type: none">•Keep out of the reach of children. Store away from direct sunlight, heat and moisture.•Discard the remainder. Do not store them.
For healthcare professional use only / /

For further information, talk to your doctor or pharmacist.

SIE①