

# Drug Information Sheet("Kusuri-no-Shiori")

Internal

Revised: 02/2016

The information on this sheet is based on approvals granted by the Japanese regulatory authority. Approval details may vary by country. Medicines have adverse reactions (risks) as well as efficacies (benefits). It is important to minimize adverse reactions and maximize efficacy. To obtain a better therapeutic response, patients should understand their medication and cooperate with the treatment.

**Brand name:** LOSARTAN POTASSIUM TABLETS 25mg 「OHARA」

**Active ingredient:** Losartan potassium

**Dosage form:** white tablet with split line, diameter 5.6mm, thick 2.9mm

**Print on wrapping:** ロサルタン K25mg 「オーハラ」, 血圧降下剤, OH-207, Losartan Potassium 25mg 「OHARA」



## Effects of this medicine

This medicine decreases blood pressure with angiotensin II-receptor antagonistic activity. It also reduces the deterioration of the renal function by reducing proteinuria to protect the kidneys.

It is usually used in the treatment of hypertension, and diabetic nephropathy in the type 2 diabetes mellitus accompanied by hypertension and proteinuria.

## Before using this medicine, be sure to tell your doctor and pharmacist

- If you have previously experienced any allergic reactions (itch, rash, etc.) to any medicines.  
If you have hyperkalemia, renal or hepatic disorder, or cerebrovascular disorder.  
If you are volume-depleted.
- If you are pregnant, possibly pregnant or breastfeeding.
- If you are taking any other medicinal products. (Some medicines may interact to enhance or diminish medicinal effects. Beware of over-the-counter medicines and dietary supplements as well as other prescription medicines.)

## Dosing schedule (How to take this medicine)

- Your dosing schedule prescribed by your doctor is<< to be written by a healthcare professional>>
- For treatment of hypertension : In general, for adults, take 1 to 2 tablets (25 to 50 mg of the active ingredient) at a time, once daily. The dosage may be adjusted according to the age or symptoms, or may be increased up to 4 tablets (100 mg) daily. Strictly follow the instructions.  
For treatment of diabetic nephropathy in the type 2 diabetes mellitus accompanied by hypertension and proteinuria : In general, for adults, take 2 tablets (50 mg of the active ingredient) at a time, once daily. The dosage may be increased up to 4 tablets (100 mg) daily according to the blood pressure level. If you have the potential to develop excessive hypotension, the initial dose is started at 1 tablet (25 mg) at a time. Strictly follow the instructions.
- If you miss a dose, take it as soon as you remember. If it is close to your next dose, do not take the missed dose. Just take the next dose at your regular time. You should never take two doses at one time.
- If you accidentally take more than your prescribed dose, consult with your doctor or pharmacist.
- Do not stop taking this medicine unless your doctor instructs you to do so.

## Precautions while taking this medicine

- The medicine may cause dizziness or lightheadedness caused by antihypertensive effects. Pay close attention to working at heights, driving a car or operating dangerous machinery.
- This medicine cannot be used by pregnant and possibly pregnant woman. If you become pregnant while taking this medicine, consult with your doctor immediately.

## Possible adverse reactions to this medicine

The most commonly reported adverse reactions include headache, dizziness, nausea, hot flush, hypotension, asthenia and fatigue. If any of these symptoms occur, consult with your doctor or pharmacist.

**The symptoms described below are rarely seen as initial symptoms of the adverse reactions indicated in brackets. If any of these symptoms occur, stop taking this medicine and see your doctor immediately.**

- hives, breathing difficulty, edema [anaphylactoid symptoms]
- swelling of the face, lips, tongue, or throat [angioedema]
- loss of appetite, general malaise, yellowing in skin and white of eyes [acute hepatitis or fulminant hepatitis]
- decreased urine output, generalized edema [renal failure]
- cold sweat, nausea, loss of consciousness [shock, faint, unconsciousness]

**The above symptoms do not describe all the adverse reactions to this medicine. Consult with your doctor or pharmacist if you notice any symptoms of concern other than those listed above.**

## Storage conditions and other information

- Keep out of the reach of children. Store away from direct sunlight, heat and moisture.

•Discard the remainder. Do not store them.

**For healthcare professional use only** / /

For further information, talk to your doctor or pharmacist.

SIE①