

# Drug Information Sheet("Kusuri-no-Shiori")

Internal

Published: 06/2018

The information on this sheet is based on approvals granted by the Japanese regulatory authority. Approval details may vary by country. Medicines have adverse reactions (risks) as well as efficacies (benefits). It is important to minimize adverse reactions and maximize efficacy. To obtain a better therapeutic response, patients should understand their medication and cooperate with the treatment.

**Brand name:**TROXIPIDE TABLETS 100mg「OHARA」

**Active ingredient:**Troxipide

**Dosage form:**white tablet, diameter: 7.1 mm, thickness: 3.5 mm

**Print on wrapping:**トロキシピド 100mg「オーハラ」, 胃炎・胃潰瘍治療剤 Troxipide 100mg「OHARA」



## Effects of this medicine

This medicine improves circulation abnormalities by increasing blood flow in the gastric mucosa, normalizes gastric mucosal component and promotes increase of prostaglandins that protect cells in the gastric mucosa, thereby improving symptoms in gastric ulcer and gastritis.

It is usually used to treat gastric ulcer and to improve gastric mucosal lesions (erosion, bleeding, redness and edema) in acute gastritis and acute exacerbation of chronic gastritis.

## Before using this medicine, be sure to tell your doctor and pharmacist

- If you have previously experienced any allergic reactions (itch, rash, etc.) to any medicines.
- If you are pregnant or breastfeeding.
- If you are taking any other medicinal products. (Some medicines may interact to enhance or diminish medicinal effects. Beware of over-the-counter medicines and dietary supplements as well as other prescription medicines.)

## Dosing schedule (How to take this medicine)

- Your dosing schedule prescribed by your doctor is(( to be written by a healthcare professional))
- In general, for adults, take 1 tablet (100 mg of the active ingredient) at a time, 3 times a day after meals. The dosage may be adjusted according the age or symptoms. Strictly follow the instructions.
- If you miss a dose, take the missed dose as soon as you remember. If it is almost time for the next dose, skip the missed dose and continue your regular dosing schedule. You should never take two doses at one time.
- If you accidentally take more than your prescribed dose, consult with your doctor or pharmacist.
- Do not stop taking this medicine unless your doctor instructs you to do so.

## Precautions while taking this medicine

### Possible adverse reactions to this medicine

The most commonly reported adverse reactions include constipation, diarrhea, nausea/vomiting, heartburn and rash/itching sensation. If any of these symptoms occur, consult with your doctor or pharmacist.

**The symptoms described below are rarely seen as initial symptoms of the adverse reactions indicated in brackets. If any of these symptoms occur, stop taking this medicine and see your doctor immediately.**

- hives, respiratory distress, dizziness on standing up/dizziness [shock, anaphylactoid symptoms]
- general malaise, loss of appetite, yellow discoloration of the skin and the white of eyes [liver dysfunction, jaundice]

**The above symptoms do not describe all the adverse reactions to this medicine. Consult with your doctor or pharmacist if you notice any symptoms of concern other than those listed above.**

### Storage conditions and other information

- Keep out of the reach of children. Store away from direct sunlight, heat and moisture.
- Discard the remainder. Do not store them.

### For healthcare professional use only / /

For further information, talk to your doctor or pharmacist.

SIE①